



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Agricultural methane emissions and the potential formitigation

**Citation for published version:**

Smith, P, Reay, D & Smith, J 2021, 'Agricultural methane emissions and the potential formitigation', *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, vol. 379, no. 2210. <<https://royalsocietypublishing.org/doi/full/10.1098/rsta.2020.0451?af=R>>

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY A

MATHEMATICAL, PHYSICAL AND ENGINEERING SCIENCES

## Agricultural Methane Emissions and the Potential for Mitigation

|   |   |
|---|---|
| Journal:  | <i>Philosophical Transactions A</i>   |
| Manuscript ID   | RSTA-2020-0451.R1   |
| Article Type:   | Research  |
| Date Submitted by the Author:   | n/a   |
| Complete List of Authors:   | Smith, Pete; University of Aberdeen, Institute of Biological & Environmental Sciences<br>Reay, Dave; University of Edinburgh<br>Smith, Jo; University of Aberdeen, Institute of Biological & Environmental Sciences |
| Issue Code (this should have already been entered and appear below the blue box, but please contact the Editorial Office if it is not present): | DM10201   |
| Subject:  | Biogeochemistry < EARTH SCIENCES, Computer modelling and simulation < COMPUTER SCIENCE, Environmental Chemistry (67) < CHEMISTRY (1002)   |
| Keywords:   | Methane, Agriculture, Emissions, Climate Change Mitigation  |
|   |   |

SCHOLARONE™  
Manuscripts

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

**Author-supplied statements**

Relevant information will appear here if provided.

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

***Ethics***

*Does your article include research that required ethical approval or permits?:*  
This article does not present research with ethical considerations

*Statement (if applicable):*  
CUST\_IF\_YES\_ETHICS :No data available.

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

***Data***

*It is a condition of publication that data, code and materials supporting your paper are made publicly available. Does your paper present new data?:*  
My paper has no data

*Statement (if applicable):*  
CUST\_IF\_YES\_DATA :No data available.

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

***Conflict of interest***

I/We declare we have no competing interests

*Statement (if applicable):*  
CUST\_STATE\_CONFLICT :No data available.

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

***Authors' contributions***

This paper has multiple authors and our individual contributions were as below

*Statement (if applicable):*  
PS wrote the first draft of the manuscript. PS, DR and JS drafted individual sections and edited / revised the manuscript drafts. All authors read and approved the manuscript.

# Agricultural Methane Emissions and the Potential for Mitigation

Pete Smith<sup>1\*</sup>, Dave Reay<sup>2</sup> & Jo Smith<sup>3</sup>

<sup>1</sup> Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen, AB24 3UU, UK. Orcid iD: 0000-0002-3784-1124

<sup>2</sup> School of Geosciences and Edinburgh Centre for Carbon Innovation, University of Edinburgh, High School Yards, Edinburgh, EH1 1LZ, UK. Orcid iD: 0000-0001-8764-3495

<sup>3</sup> Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen, AB24 3UU, UK. Orcid iD: 0000-0001-6984-6766

**Keywords:** Methane, agriculture, mitigation, greenhouse gas, emissions

## Summary

Agriculture is the largest anthropogenic source of methane (CH<sub>4</sub>), emitting 145 Tg CH<sub>4</sub> y<sup>-1</sup> to the atmosphere in 2017. The main sources are enteric fermentation, manure management, rice cultivation and residue burning. There is significant potential to reduce CH<sub>4</sub> from these sources, with bottom-up mitigation potentials of ~10.6, 10, 2 and 1 Tg CH<sub>4</sub> y<sup>-1</sup> from rice management, enteric fermentation, manure management and residue burning. Other system-wide studies have assumed even higher potentials of 4.8 to 47.2 Tg CH<sub>4</sub> y<sup>-1</sup> from reduced enteric fermentation, and 4 to 36 Tg CH<sub>4</sub> y<sup>-1</sup> from improved rice management. Biogas (a methane-rich gas mixture generated from anaerobic decomposition of organic matter and used for energy) also has potential to reduce unabated CH<sub>4</sub> emissions from animal manures and human waste. In addition to these supply-side measures, interventions on the demand-side (shift to a plant-based diet and a reduction in total food loss and waste by 2050) would also significantly reduce methane emissions, perhaps in the order of >50 Tg CH<sub>4</sub> y<sup>-1</sup>. While there is an pressing need to reduce emissions of long-lived greenhouse gases (CO<sub>2</sub> and N<sub>2</sub>O) due to their persistence in the atmosphere, despite CH<sub>4</sub> being a short-lived greenhouse gas, the urgency of reducing warming means we must reduce any GHG emissions we can as soon as possible. Because of this, mitigation actions should focus on reducing emissions of all the three main anthropogenic greenhouse gases, including CH<sub>4</sub>.

\*Author for correspondence (pete.smith@abdn.ac.uk).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

---

# 1 Emissions of methane from agriculture

The main sources of methane (CH<sub>4</sub>) emissions from agriculture are enteric fermentation, manure management, rice cultivation and residue burning, with FAOSTAT being the main source of statistics on agricultural emissions [1].

Enteric CH<sub>4</sub> is produced under anaerobic conditions by a diverse community of methanogenic archaea, using mainly hydrogen and CO<sub>2</sub> as substrates, although smaller amounts are produced using formate and methyl compounds as alternatives to hydrogen [2]. The quantity of feed consumed by a ruminant largely determines the quantity of CH<sub>4</sub> emitted, though the type and quality of the animal feed also influence emissions [3,4]. The species of ruminant, an individual's digestive physiology and the makeup of the resident microbial population can also influence the quantity of CH<sub>4</sub> it produces [5,6,7].

Methane production from animal wastes is also an anaerobic microbial process and occurs mostly when animal wastes are stored (manure management). Smaller quantities are produced from wastes deposited directly onto the ground. Manure type (e.g. wet versus dry), storage method, storage duration, manure chemical composition and temperature all influence the quantity of methane produced per unit of substrate [8].

Methane emissions from paddy rice occur when soils are flooded, which creates anaerobic conditions suitable for methanogenic microorganisms to produce CH<sub>4</sub>. While methanotrophs are able to oxidise some of the CH<sub>4</sub> produced, there is still a large net emission from paddy rice fields [9]. Global cropland CH<sub>4</sub> emissions are dominated by rice production, with 90% of emissions from tropical Asia, more than half from China and India combined [10], and a small contribution to the global CH<sub>4</sub> soil sink from other croplands (see section 1.2).

Residue burning releases CH<sub>4</sub> through incomplete combustion of biomass, though the quantity is small compared to enteric fermentation, manure management and rice cultivation [1].

## 1.1 The global methane budget and the contribution of agriculture

Global CH<sub>4</sub> emissions were 596 (572–614) Tg y<sup>-1</sup>, partly offset by a CH<sub>4</sub> sink of 571 (540–585) Tg y<sup>-1</sup> in 2017 (see section 1.2) [11]. Of total CH<sub>4</sub> emissions, bottom-up and top-down estimates of the anthropogenic component were 380 (359–407) and 364 (340–381) Tg y<sup>-1</sup>, respectively in 2017 [11]. Of total anthropogenic CH<sub>4</sub> emissions, the majority were attributable to the agriculture and waste sector, with bottom-up and top-down estimates of 213 (198–232) and 227 (205–246) Tg y<sup>-1</sup>, respectively in 2017 [11], with bottom-up estimates of emissions suggesting that 68% of these are from agriculture (Figure 1).

[Figure 1 here]

Within the agricultural sector, enteric fermentation and manure management together contributed 115 (110–121) Tg CH<sub>4</sub> y<sup>-1</sup>, rice cultivation contributed 30 (24–40) Tg CH<sub>4</sub> y<sup>-1</sup>, with the remainder from landfills and waste (68 [64–71] Tg CH<sub>4</sub> y<sup>-1</sup>) in 2017 [11]. Enteric fermentation represents about 30–32% of total anthropogenic CH<sub>4</sub> emissions. Enteric fermentation is responsible for about 90% of all livestock derived CH<sub>4</sub> emissions, with cattle (77%) being the dominant source [12]. Manure management emissions are dominated by pigs (~42%) and cattle (~41%) [12].

Additional managed land-based emission sources in 2017, though not accounted for in the agriculture sector, were 16 (11–24) and 13 (10–14) Tg CH<sub>4</sub> y<sup>-1</sup>, for biomass burning and biofuel burning, respectively [11]. Agricultural CH<sub>4</sub> emissions in 2017 have increased since the early 2000s (2000–2006) by 12.7% for enteric fermentation and manure management, and 7.1% for rice cultivation [11]. Changes in agricultural CH<sub>4</sub> emissions for 1961 to 2017 are shown in figure 2.

[Figure 2 here]

Regionally, for enteric fermentation emissions, largest emissions are found in Asia followed by Latin America, OECD-90, Africa and the Middle East and Economies in transition [10, 13]. For manure management, largest emissions are seen in OECD-90 and Asia, and for rice emissions, Asia has larger emissions than all other world regions together [10]. The increase in agricultural CH<sub>4</sub> emissions from the early 2000s and 2017 was largely seen in South America, Africa (7–9 Tg y<sup>-1</sup>) - largely from enteric fermentation and manure, and South

*Phil. Trans. R. Soc. A.*

Asia/Oceania (of 9-10 Tg y<sup>-1</sup>) – from paddy rice, enteric fermentation and manure. Estimates of agricultural emissions of CH<sub>4</sub> in Europe fell by -1.4 to -2.8 Tg y<sup>-1</sup>, for top-down and bottom-up methods, respectively [11].

### 1.2 Sinks of methane in agriculture

As noted in Section 1.1 above, there are large natural sinks for CH<sub>4</sub>. Most of the CH<sub>4</sub> sink is in the atmosphere, which includes reaction with tropospheric hydroxy (OH) radicals to produce carbon dioxide (CO<sub>2</sub>) and water, and chlorine (Cl) radicals in the troposphere and the stratosphere. The other significant sink, estimated to be responsible for uptake of 30 (11-49) or 40 (37-47) Tg CH<sub>4</sub> y<sup>-1</sup> in 2017 from bottom-up and top-down measurements, respectively, is the soil [11].

Cultivation of land for agriculture can significantly reduce the sink capacity of soils to oxidize CH<sub>4</sub> [14]. Mineral soils under forests and other natural vegetation act as the strongest CH<sub>4</sub> sink, followed by grasslands, with the sink strength weakest in cultivated soils and those receiving nitrogen fertilizer [7,14,15]; as such, as cropland has expanded, the CH<sub>4</sub> sink strength of soils globally will have declined [14]. When mineral soils become anaerobic, the net flux to the atmosphere can be positive, with waterlogged soils becoming a CH<sub>4</sub> source, often with large emission rates [16]. When soils are deliberately flooded, e.g. for paddy rice cultivation, they can become very large global sources of CH<sub>4</sub> as described in section 1.1 [7].

### 1.3 Metrics of the climate warming effect of methane

For comparability with other greenhouse gases, the radiative forcing of CH<sub>4</sub> is often expressed in terms of CO<sub>2</sub> equivalents, calculated using a global warming potential (GWP) over a 100-year time horizon (GWP<sub>100</sub>). National greenhouse gas inventories, to date, have used a GWP<sub>100</sub> value of 25 from the IPCC Fourth Assessment Report (1 kg of CH<sub>4</sub> is equivalent to 25 kg of CO<sub>2</sub>). The GWP<sub>100</sub> of CH<sub>4</sub> has frequently been updated as scientific understanding has improved and was quoted as 21, 25 and 28 in the IPCC 2<sup>nd</sup>, 4<sup>th</sup> and 5<sup>th</sup> Assessment Reports, respectively. When feedbacks are included, the GWP<sub>100</sub> value for CH<sub>4</sub> was estimated to be 34 in the IPCC 5<sup>th</sup> Assessment Report.

Given the relatively short atmospheric lifetime of CH<sub>4</sub> – 12.4 years compared to 121 for nitrous oxide and 300- >1000 years for CO<sub>2</sub> – some have argued that GWP<sub>100</sub> is not a useful metric for assessing the contribution of

CH<sub>4</sub> to climate warming [17]. Instead, they propose a metric that reports equivalent emissions, based on whether sustained changes in the emission rates of short-lived gases (like CH<sub>4</sub>) would result in a similar warming contribution to an individual, one-off CO<sub>2</sub> emission [18, 19]. This metric is known as GWP\* [17,18,19].

The consequences of using the GWP<sub>100</sub> and GWP\* metrics for assessing the climate warming caused by CH<sub>4</sub> are very different. Instead of providing a snapshot of CH<sub>4</sub> emissions at a single point in time, the calculation underpinning GWP\* expresses the warming impacts of changes in the *rate of emissions of CH<sub>4</sub>* as equivalent to a large pulse emission of CO<sub>2</sub> [20]. Using GWP\* as the metric for CH<sub>4</sub>, if CH<sub>4</sub> emissions remain constant there is no additional warming, unlike for CO<sub>2</sub> (or other long-lived gases) where each additional tonne of CO<sub>2</sub> added to the atmosphere causes additional warming. This has led some sectors of the agricultural industry, particularly in the livestock sector, to make statements such as: “This means that the CH<sub>4</sub> emissions of a herd of 100 cows today are simply replacing the emissions that were first produced when that herd was established by a previous generation of farmers. There was an initial pulse of warming when the herd was established, but there is no ongoing warming from that herd” [21]. These statements are used to support arguments that grazed livestock are part of the climate solution [21]. The assertion has been challenged [22] and the authors of the GWP\* themselves note, “while some ongoing CH<sub>4</sub> emissions may be able to give no further temperature increases from those emissions, maintaining these emissions into the future means they will continue to contribute to our elevated temperatures, and the resulting climate damages we will experience” [20]. It is a fundamental, metric-independent reality that emitting less methane will mean having a smaller impact on the climate.

## 2 Reducing methane emissions from agriculture

As outlined in section 1.1, the main sources of CH<sub>4</sub> emissions from agriculture are from rice production, enteric fermentation, manure management and residue burning. The technical options for reducing emissions from these sources are described below, along with their estimated global mitigation potential, summarised in Figure 3.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 2.1 Mitigation opportunities in rice production

Changes in rice management have the potential to significantly decrease paddy rice soil CH<sub>4</sub> emissions [10,23]. Mid-season drainage is the main mitigation option with other mitigation measures including changed fertilizer practices and tillage/residue management [10].

Emissions during the growing season can be reduced by many practices [24,25,26]. Mid-season drainage effectively reduces CH<sub>4</sub> emissions [27,28], although this benefit may be partly offset by higher nitrous oxide emissions, and the practice may be constrained by water supply. Mid-season drainage is now becoming prevalent in many rice-growing areas [7]. Rice cultivars with low exudation rates could offer an important CH<sub>4</sub> mitigation option [26]. In the off-rice season, CH<sub>4</sub> emissions can be reduced by improved water management, especially by keeping the soil as dry as possible and avoiding waterlogging [29,30,31,32]. Methane emissions can also be reduced by adjusting the timing of organic residue additions (e.g. incorporating organic materials in the dry period rather than in flooded periods [33,34]) and composting the residues before incorporation.

The estimated global mitigation potential for rice management has been estimated to be ~8, 9 and 10 Tg CH<sub>4</sub> y<sup>-1</sup> at carbon prices of 20, 50 and 100 US\$ tCO<sub>2</sub>e, respectively [10,23].

## 2.2 Mitigation opportunities for enteric fermentation

Practices for reducing enteric CH<sub>4</sub> emissions fall into three general categories: a) improved feeding practices, b) use of specific agents or dietary additives, and c) longer term management changes and animal breeding. Additional options to reduce emissions arise from reducing ruminant livestock numbers, enabled by demand-side changes (dietary change and reduced food loss/waste; discussed further in section 3.1).

For improved feeding practices, CH<sub>4</sub> emissions can be reduced by feeding livestock more concentrates which normally replace forage [35,36,37,38]. Although concentrates may increase daily CH<sub>4</sub> emissions, emissions per unit of feed intake and per unit product (emission intensity) are almost always reduced. The net benefit, however, depends on reduced animal numbers or younger age at slaughter for beef animals and on how the practice affects emissions when producing and transporting the concentrates [39,40]. Other practices that can

reduce enteric CH<sub>4</sub> emissions include adding oils to the diet [41,42] and improving pasture quality, especially in less developed regions, because it improves animal productivity and reduces the proportion of energy lost as CH<sub>4</sub> [43,44,45].

A wide range of specific agents and dietary additives have been tested, mostly aimed at suppressing methanogenesis. These include ionophores, which are antibiotics that can reduce CH<sub>4</sub> emissions [46,47,48], but their effect may be transitory [49] and they have been banned in some jurisdictions, such as the European Union. Halogenated compounds which inhibit methanogenic bacteria [50,51] have also been tested, but their effects, too, are often transitory and they can have side effects such as reduced calorie intake. Probiotics, such as yeast culture, have shown only small, insignificant effects [48], but selecting strains specifically for CH<sub>4</sub> reducing ability could improve results [52]. Propionate precursors, such as fumarate or malate, reduce CH<sub>4</sub> formation by acting as alternative hydrogen acceptors [53], but are effective only at high doses and are therefore expensive [54]. Vaccines against methanogenic bacteria have been developed but are not yet commercially available [55]. Bovine somatotrophin (bST) and hormonal growth implants do not specifically suppress CH<sub>4</sub> formation, but by improving animal performance [56,57] they can reduce the emission intensity (emissions per unit of product) of meat/dairy [58,59], but like ionophores, are banned in some jurisdictions, such as the European Union. Some natural feed additives, such as seaweed, have been tried [60]

Longer term management changes and animal breeding includes increasing productivity through breeding and better management practices, which spreads the energy cost of maintenance across a greater feed intake, often reducing CH<sub>4</sub> output per unit of animal product [61]. With improved efficiency, meat-producing animals reach slaughter weight at a younger age, with reduced lifetime emissions [62]. The whole system effects of such practices are not clear, however; for example, selecting for higher yield might reduce fertility, requiring more replacement animals [40].

Mitigation potential from reducing enteric fermentation from livestock has been estimated to be ~6.4, 8.5 and 10.6 Tg CH<sub>4</sub> y<sup>-1</sup> at carbon prices of 20, 50 and 100 US\$ tCO<sub>2</sub>e, respectively [10,23].

## 2.3 Mitigation opportunities in manure management

*Phil. Trans. R. Soc. A.*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

---

Animal manures can release significant amounts of CH<sub>4</sub> during storage, but the magnitude of these emissions varies. Methane emissions from manure stored in lagoons or tanks can be reduced by cooling or covering the sources, or by capturing the CH<sub>4</sub> emitted [63,64,65,66]. The manures can also be digested anaerobically to maximize retrieval of CH<sub>4</sub> as an energy source [63,67]; see section 2.5).

Storing and handling the manures in solid, rather than liquid form, can suppress CH<sub>4</sub> emissions but may increase nitrous oxide formation [66]. For most livestock production systems globally, there is limited opportunity for manure management, as treatment or storage - excretion happens in the field and handling for fuel or fertility amendment occurs when it is dry and CH<sub>4</sub> emissions are negligible [66]. Emissions from manure might be curtailed somewhat by altering feeding practices [69] or by composting the manure [70], but these mechanisms and the system-wide impacts have not been widely explored.

Mitigation potential from improved manure management has been estimated to be ~0.4, 1 and 2 Tg CH<sub>4</sub> y<sup>-1</sup> at carbon prices of 20, 50 and 100 US\$ tCO<sub>2</sub>e, respectively [10,23].

## 2.4 Mitigation opportunities for residue burning

Strategies to reduce residue burning are often promoted to improve air quality and address a mix of long- and short-lived climate pollutants [7]. Since residue burning is responsible for just over 1 Tg CH<sub>4</sub> y<sup>-1</sup> (Figure 2, [1]), the total cessation of crop residue burning would have a maximum mitigation potential of ~1 Tg CH<sub>4</sub> y<sup>-1</sup>. Figure 3 summarises the mitigation potentials described in this section.

## 2.5 Potential for biogas

Biogas is a methane-rich gas mixture, generated from anaerobic decomposition of organic matter that can be burnt to release energy. Use of organic wastes in production of biogas has potential to change net CH<sub>4</sub> emissions in two ways. Collection of organic wastes for use as a feedstock for biogas production may reduce CH<sub>4</sub> emissions by removing wastes from the environment where uncontrolled anaerobic decomposition can result in significant emissions of CH<sub>4</sub> [8]. However, emissions of CH<sub>4</sub> may also be increased by CH<sub>4</sub> leakage from the biogas digesters [71], piping [2] and appliances [73,4]. The net effect on CH<sub>4</sub> emissions is a balance between these different processes. While we focus here only on the impact of biogas on CH<sub>4</sub> emissions, it

should be noted that using organic wastes in biogas production has further impacts on total greenhouse gas emissions by potentially replacing fossil fuels [75], reducing deforestation associated with use of wood as a fuel [77,77] and increasing soil carbon sequestration associated with application of bioslurry as an organic fertilizer [78]. These latter impacts are not discussed further here.

*Reduced emissions of methane from organic wastes* – The global emissions of CH<sub>4</sub> from deposited and stored manures is estimated to be 9.9 Tg y<sup>-1</sup> (Figure 2; [1]). The maximum potential reduction in CH<sub>4</sub> emissions associated with prevention of uncontrolled anaerobic decomposition of manures is 9.9 Tg y<sup>-1</sup>, but since only stored manures can be used for biogas production, this maximum potential is likely to be substantially lower. In practice, anaerobic digestion can only be implemented in locations with sufficient access to water [79]. Consistency of supply of water is also important, with seasonal breaks in supply likely to increase the proportion of digesters that are abandoned [80]. Therefore, the actual potential for reduction in CH<sub>4</sub> emissions using anaerobic digestion is likely to be significantly less than this maximum potential. The proportion of the CH<sub>4</sub> produced that leaks from the digester, pipes and biogas appliances is dependent on the scale of the system and the sophistication of the technology used; large-scale, state-of-the-art plants are likely to leak a much lower proportion of CH<sub>4</sub> produced than simple, small-scale systems. Net emissions also depend on the counterfactual emissions from the energy that is replaced by biogas; for example, for household cookstoves, emissions of CH<sub>4</sub> during combustion are 57 mg per MJ energy delivered for biogas, compared to 8.9 mg MJ<sup>-1</sup> for LPG, 600 mg MJ<sup>-1</sup> for wood, 1300 mg MJ<sup>-1</sup> for coal and 7100 mg MJ<sup>-1</sup> for dung. Therefore, combustion losses of CH<sub>4</sub> from cookstoves are increased only compared to LPG, whereas by comparison to wood, coal and dung, combustion losses are very much reduced [71].

Leaks of CH<sub>4</sub> from biogas digesters can occur from any openings in the digester tank; for example, in fixed dome digesters, the inlet and outlet are open to the atmosphere, so any CH<sub>4</sub> produced in these locations can be lost, while in floating drum digesters, any CH<sub>4</sub> produced from the small volume of manure on the outside of the upper drum can be lost. These losses from well-maintained small-scale digesters in India have been estimated to be 14 – 17% of the CH<sub>4</sub> produced in fixed dome digesters [81], and 5 – 8% in floating drum digesters [82]. Cracks in the digester body or gas tubing due to poor maintenance can result in further unintentional losses of CH<sub>4</sub>. Even in well-maintained large-scale agricultural digesters in Canada, these losses

*Phil. Trans. R. Soc. A.*

were estimated to average 3.1% of the CH<sub>4</sub> produced [83], whereas in less well-maintained systems in China, fugitive losses due to poor maintenance were estimated to be as high as 10% [71]. However, the largest source of CH<sub>4</sub> emissions from biogas digesters may be due to the intentional venting (without flaring) of excess biogas; these losses were estimated in a study of small scale digesters in Thailand to be 15% of the CH<sub>4</sub> produced [84], and in southern Vietnam to be as high as 36.6% [72]. In larger scale systems, alternative uses are usually found for excess biogas, and any further excess is usually converted to CO<sub>2</sub> by flaring. Bruun et al. [71] estimated that typical total CH<sub>4</sub> losses due to leaks and venting from small-scale biogas digesters is in the region of 40% of the CH<sub>4</sub> produced, and estimated that in 2014, this amounted to a global total of ~4.5 Tg y<sup>-1</sup>.

Therefore, while anaerobic digestion has potential to reduce CH<sub>4</sub> emissions from uncontrolled anaerobic decomposition of manures by up to 9.9 Tg y<sup>-1</sup>, losses due to leaks from digesters, pipes and appliances are likely to be in the region of ~4.5 Tg y<sup>-1</sup>. Therefore, the net potential impact of anaerobic digestion on CH<sub>4</sub> emissions could be to increase CH<sub>4</sub> emissions by up to 4.5 Tg y<sup>-1</sup> if no reduction in uncontrolled decomposition is achieved, or to reduce CH<sub>4</sub> emissions by up to 4.4 Tg y<sup>-1</sup>. Future initiatives to increase implementation of anaerobic digestion must therefore be combined with improvements in maintenance of digesters in order to achieve maximum benefits in CH<sub>4</sub> emission reduction and avoid increased emissions.

[Figure 3 here]

### 3 Reducing methane emissions from the food system

#### 3.1 Dietary change

Food supply chain and demand-side interventions that save CH<sub>4</sub> emissions at the production phase, such as reduced supply chain loss and waste, also have an important role to play in this sector [85]. Since enteric fermentation dominates agricultural CH<sub>4</sub> emissions, any transition away from ruminant livestock will reduce CH<sub>4</sub> emissions [86].

On an emissions intensity basis (greenhouse gas emissions per unit mass, protein or energy), numerous studies have shown the climate impact of ruminant meat to be 10-100 times greater than plant-based foods [86,

87,88,89,90], so a shift away from ruminant meat and dairy, toward plant-based products in the diet, greatly reduces the climate footprint of food, largely by reducing CH<sub>4</sub> emissions [89,90,91,92,93,94,94].

A shift toward meat from monogastrics (e.g. pig and poultry) also lowers CH<sub>4</sub> emissions, since they produce no enteric CH<sub>4</sub>, although emissions from manure management remain. In a meta-analysis, Aleksandrowicz et al. [94] showed that a vegan diet reduced emissions by 45% (>20->70%) relative to current average diets, vegetarian diet reduced emissions by ~30% (15%~60%) while transition to meat from monogastrics reduced emissions by ~20% (~5%~35%). It is worth noting that dietary transitions for individuals do not have to be absolute; any reduction in ruminant product consumption will reduce CH<sub>4</sub> emissions associated with diets. A dietary transition to one in which every person on the planet eats according to healthy dietary guidelines would deliver significant CH<sub>4</sub> emission reductions [93].

Since CH<sub>4</sub> from enteric fermentation are around 100 Tg CH<sub>4</sub> y<sup>-1</sup> in 2017 (see Figure 2; [1]), the maximum technical emission reduction potential (with no ruminant meat or dairy consumption) would be 100 Tg CH<sub>4</sub> y<sup>-1</sup>, but Roe et al. [95] model an equivalent of 50% of the human population, which would halve CH<sub>4</sub> emissions from enteric fermentation, ceasing eating meat and dairy to deliver a land / food system that is compliant with a 1.5°C world (see section 4.2).

### 3.2 Waste reduction

Food supply chain and demand-side interventions that save CH<sub>4</sub> emissions at the production phase, such as reduced supply chain loss and waste, also have an important role to play in this sector. An estimated 26% of food produced globally is lost or wasted each year, equivalent to 6% of global anthropogenic greenhouse gas emissions [89]. Methane-intensive foods, such as ruminant meat and dairy, play a disproportionately large role in these food wastage emissions and one that has continued to expand over the past half century [85].

In developed nations, the bulk of these losses occur in the consumer phase, with avoidable wastage of milk in UK households, for example, being estimated at 290 thousand tonnes each year [96]. Applying a simplistic global average CH<sub>4</sub> emission factor (48kg CH<sub>4</sub> per tonne of milk [based on [97] and assuming CH<sub>4</sub> comprises

*Phil. Trans. R. Soc. A.*

50% of global average footprint of 2.4kg CO<sub>2</sub>e per kg fat and protein-corrected milk and GWP<sub>100</sub> of 25)) to these consumer-phase milk losses equates to around 14 thousand tonnes of CH<sub>4</sub> emission per year for the UK alone.

Similarly, for dairy milk in the US, the huge volumes wasted represent a very large CH<sub>4</sub> emissions penalty, but by reducing losses in both the retail and consumer phases, Thoma et al. [98] estimate that emissions from US milk could be reduced by 23%. As such, reducing food loss and waste represents a potentially powerful, albeit indirect, CH<sub>4</sub> mitigation strategy for global agriculture.

## 4 Future prospects

### 4.1 Climate change impacts on future agricultural methane fluxes

Climate change itself may alter future CH<sub>4</sub> fluxes from agriculture, and so the efficacy of mitigation measures. For the livestock sector, changes in feed quantity and quality, increased animal heat stress and manure fermentation rates, and increased pest and disease impacts, may all serve to enhance emissions [8]. The net effect globally remains highly uncertain, with wide variation in impacts likely between different regions and production systems. Adaptation will play a central role here in terms of buffering the impacts of climate change at local scales, such as through use of shading, ventilation and livestock management strategies in the case of extreme heat events [99].

For cropland systems the projected impacts of climate change on CH<sub>4</sub> fluxes are relatively minor and largely stem from changes in soil moisture, such as drying of waterlogged mineral soils reducing methanogenesis. More important will be the effects on CH<sub>4</sub> emissions from rice agriculture. Here, reduced soil moisture may substantially reduce emissions in some rain-fed systems [100], while in irrigated systems higher temperatures combined with enhanced atmospheric CO<sub>2</sub> concentrations can greatly increase emissions [101]. As with the livestock sector, variation across locations and production systems will be large.

Overall, the greatest impacts of climate change on CH<sub>4</sub> emissions from agriculture are likely to arise indirectly through effects on production efficiency. While this includes heat stress, drought, disease and other ‘on-farm’



impacts, it is also relevant right along the food supply chain, for instance, higher temperatures increasing food spoilage rates. As discussed earlier, given the current magnitude of loss and wastage of CH<sub>4</sub>-intensive foods, such as milk, any climate change impacts that exacerbate these losses risk an upstream ripple effect of increased on-farm emissions.

The net impact of such climate change- CH<sub>4</sub> feedbacks on emissions from agriculture at a global scale is likely to be dwarfed by future changes in food demand, land use, and food system management practices (including those focussed on mitigation). Nevertheless, CH<sub>4</sub> mitigation strategies in agriculture must be cognisant of these feedbacks, make the most of any synergies with climate adaptation and avoid any undermining of food system resilience.

## 4.2 Methane reduction in climate stabilization pathways

While some studies have suggested that future temperature targets could be achieved without major reductions in ruminant/agricultural methane emissions (102,103,104), Roe et al. [95], synthesising previous top-down and bottom-up estimates of mitigation in agriculture propose a 25% reduction in agricultural non-CO<sub>2</sub> emissions by 2050, compared to business as usual, in their implementation roadmap for the land sector. Priority regions for reducing CH<sub>4</sub> emissions from enteric fermentation and manure management are China, India, Brazil, EU, US, Australia, Russia and Latin America (Brazil, Argentina, Mexico, Colombia, Paraguay, Bolivia). Priority regions for reducing CH<sub>4</sub> emissions by improving water and residue management of rice fields, and manure management are in Asia, namely India, China, Indonesia, Thailand, Bangladesh, Vietnam, Philippines. Globally, this translates to mitigation from reduced enteric fermentation from better feed and animal management of 4.8 to 47.2 Tg CH<sub>4</sub> y<sup>-1</sup>, and 4 to 36 Tg CH<sub>4</sub> y<sup>-1</sup> from improved rice management. Note that the higher numbers in the range are somewhat higher than the potentials reported in section 2.

In their implementation roadmap for the land sector, Roe et al. [95] propose 50% of the global population shift to a plant-based diet by 2050 and a 50% reduction in total food loss and waste by 2050 compared to BAU. The priority regions for a shift to plant-based diets are developed and emerging countries, i.e. US, EU, China, Brazil, Argentina, Russia and Australia), while the priority regions for reduced food waste are China, Europe, North America and Latin America, and for reduced food loss are Southeast Asia and Sub-Saharan Africa. The

*Phil. Trans. R. Soc. A.*



estimated mitigation potential of these measures, excluding land-use change benefits, is 0.9 Gt CO<sub>2</sub>e y<sup>-1</sup> for 50% shift to plant-based diets by 2050 and 0.9 Gt CO<sub>2</sub>e y<sup>-1</sup> for a 50% reduction in food loss and waste by 2050 [95]. Not all of this, however, is through CH<sub>4</sub> reduction, and the figures include reduction in nitrous oxide emissions [105]. If 50% of the global population shift to a plant-based diet by 2050 and a 50% reduction in total food loss and waste led to a halving of enteric fermentation and manure production, the mitigation potential could be in the order of >50 Tg CH<sub>4</sub> y<sup>-1</sup>.

Methane abatement is clearly an important component of a land sector that helps to deliver a 1.5 °C world, with interventions both on the supply side (reduction in emissions from enteric fermentation, rice and manure) and the demand side (dietary shifts toward plant-based diets and reduction in food loss and waste) necessary to achieve a land sector that is compliant with the Paris Climate Agreement [95], with the IPCC in the Special Report on 1.5°C target suggesting that agricultural methane emissions need to be 24-47% below 2010 emissions in 2050 [106].

## 5 Concluding remarks

Agriculture is the largest anthropogenic source of methane, emitting 145 Tg CH<sub>4</sub> y<sup>-1</sup> to the atmosphere in 2017. The main sources are enteric fermentation, manure management, rice cultivation and residue burning. There is significant potential to reduce CH<sub>4</sub> from these sources, with mitigation potentials of ~10.6, 10, 2 and 1 Tg CH<sub>4</sub> y<sup>-1</sup> from rice management, enteric fermentation, manure management and residue burning, respectively (Figure 3). Other studies assume even higher potentials of 4.8 to 47.2 Tg CH<sub>4</sub> y<sup>-1</sup> from reduced enteric fermentation, and 4 to 36 Tg CH<sub>4</sub> y<sup>-1</sup> from improved rice management [95]. Biogas also has potential to reduce unabated CH<sub>4</sub> emissions from animal manures and human waste. In addition to these supply-side measures, interventions on the demand-side (50% of the global population shift to a plant-based diet by 2050 and a 50% reduction in total food loss and waste) would also significantly reduce methane emissions, perhaps on the order of >50 Tg CH<sub>4</sub> y<sup>-1</sup>.

While there is an pressing need to reduce emissions of long-lived greenhouse gases (CO<sub>2</sub> and N<sub>2</sub>O) due to their persistence in the atmosphere, despite CH<sub>4</sub> being a short-lived greenhouse gas, the urgency of reducing warming means we must reduce any GHG emissions we can as soon as possible. Mitigation actions should focus on reducing emissions of all the three main anthropogenic greenhouse gases, including CH<sub>4</sub>.

## Additional Information

### Ethics

There are no ethical considerations in the paper.

### Data Accessibility

This paper does not report primary data. All publicly available data sources are given. There are no Supplementary Materials.

### Authors' Contributions

PS wrote the first draft of the manuscript. PS, DR and JS drafted individual sections and edited / revised the manuscript drafts. All authors read and approved the manuscript.

### Competing Interests

The authors declare that they have no competing interests.

### Funding Statement

The input of PS contributes to projects DEVIL [NE/M021327/1] and Soils-R-GRREAT [NE/P019455/1].

## Reference

1. FAOSTAT. 2020 <http://www.fao.org/faostat/en/#data/GT> (last accessed 26th October 2020)
2. Tapio I, Snelling TJ, Strozzi F, Wallace RJ. 2017 The ruminal microbiome associated with methane emissions from ruminant livestock. *J. Anim. Sci. Biotechnol.* **8**, 7 (doi:10.1186/s40104-017-0141-0. eCollection 2017)
3. Hristov AN *et al.* 2013 *Mitigation of greenhouse gas emissions in livestock production: a review of technical options for non-CO<sub>2</sub> emissions* (eds. PJ Gerber, B Henderson, HPS Makkar). FAO Animal Prod. Health Pap. 177, Rome
4. Negussie E *et al.* 2017 Large-scale indirect measurements for enteric methane emissions in dairy cattle: a review of proxies and their potential for use in management and breeding decisions. *J. Dairy Sci.* **100**, 2433–2453 (doi:10.3168/jds.2016-12030)

5. Goopy JP *et al.* 2014 Low-methane yield sheep have smaller rumens and shorter rumen retention time. *Br. J. Nutr.* **111**, 578–585 (doi:10.1017/S0007114513002936)
6. Pinares-Patiño C, Ebrahimi SH, McEwan J, Dodds K, Clark H, Luo D. 2011 Is rumen retention time implicated in sheep differences in methane emission. *Proc. N. Z. Soc. Anim. Prod.* **71**, 219–222.
7. Shi W *et al.* 2014. Methane yield phenotypes linked to differential gene expression in the sheep rumen microbiome. *Genome Res.* **24**, 1517–1525 (doi:10.1101/gr.168245.113)
8. Reay DS, Smith P, Christensen, TR, James RH, Clark H. 2018. Methane and global environmental change. *Annu Rev Environ Resour* **43**, 165-192 (doi:10.1146/annurev-environ-102017-030154)
9. Yan X, Akiyama H, Yagi K, Akimoto H. 2009 Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Intergovernmental Panel on Climate Change Guidelines. *Glob. Biogeochem. Cycles* **23**, GB2002 (doi:10.1029/2008GB003299)
10. Smith P *et al.* 2014 Agriculture, forestry and other land use (AFOLU). In *Climate change 2014: mitigation of climate change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds O Edenhofer *et al.*), pp. 811-922. Cambridge, UK: Cambridge University Press.
11. Jackson, RB *et al.* 2020 Increasing anthropogenic methane emissions arise equally from agricultural and fossil fuel sources. *Environ. Res. Lett.* **15**, 071002 (doi:10.1088/1748-9326/ab9ed2).
12. Gerber PJ, *et al.* 2013 *Tackling Climate Change Through Livestock: A Global Assessment of Emissions and Mitigation Opportunities*. Rome: Food Agric. Org. U.N.
13. Smith P *et al.* 2013 How much land-based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? *Glob. Change Biol.* **19**, 2285-2302 (doi:10.1111/gcb.12160)
14. Tate KR. 2015 Soil methane oxidation and land-use change—from process to mitigation. *Soil Biol. Biochem.* **80**, 260–272 (doi:10.1016/j.soilbio.2014.10.010)
15. Dutaur L, Verchot LV. 2007 A global inventory of the soil CH<sub>4</sub> sink. *Glob. Biogeochem. Cycles* **21**, GB4013 (doi:10.1029/2006GB002734)
16. Le Mer J, Roger P. 2001 Production, oxidation, emission and consumption of methane by soils: a review. *Eur. J. Soil Biol.* **37**, 25–50 (doi:10.1016/S1164-5563(01)01067-6)

17. Allen MR *et al.* 2016 New use of global warming potentials to compare cumulative and short-lived climate pollutants. *Nat. Clim. Change* **6**, 773 (doi:10.1038/nclimate2998)
18. Cain M *et al.* 2019 Improved calculation of warming-equivalent emissions for short-lived climate pollutants. *NPJ Clim. Atmos. Sci.* **2**, 29 (doi:10.1038/s41612-019-0086-4)
19. Lynch J, Cain M, Pierrehumbert R, Allen M. 2020 Demonstrating GWP\*: a means of reporting warming-equivalent emissions that captures the contrasting impacts of short- and long-lived climate pollutants. *Environ. Res. Lett.* **15**, 044023 (doi:10.1088/1748-9326/ab6d7e)
20. Lynch J, Garnett T, Persson M, Rööös E, Reisinger A 2020 *Methane and the sustainability of ruminant livestock* (Foodsource: building blocks). Food Climate Research Network, University of Oxford.
21. Costain f. 2019 Livestock are not the global warming enemy. *Vet. Rec.* **185**, 449 (doi:10.1136/vr.l5963)
22. Smith P, Balmford A 2020 Climate change: 'no get out of jail free card'. *Vet. Rec.* **186**, 71. (doi:10.1136/vr.m190)
23. Smith P *et al.* 2008. Greenhouse gas mitigation in agriculture. *Phil. Trans. R. Soc. B.* **363**, 789-813. (doi:10.1098/rstb.2007.2184)
24. Yagi K, Tsuruta H, Minami K 1997 Possible options for mitigating methane emission from rice cultivation. *Nutr. Cycl. Agroecosyst.* **49**, 213–220 (doi:10.1023/A:1009743909716)
25. Wassmann R, Lantin RS, Neue HU, Buendia LV, Corton TM, Lu Y 2000 Characterization of methane emissions from rice fields in Asia. III. Mitigation options and future research needs. *Nutr. Cycl. Agroecosyst.* **58**, 23–36 (doi:10.1023/A:1009874014903)
26. Aulakh MS, Wassmann R, Bueno C, Rennenberg H. 2001 Impact of root exudates of different cultivars and plant development stages of rice (*Oryza sativa* L.) on methane production in a paddy soil. *Plant Soil* **230**, 77–86 (doi:10.1023/A:1004817212321)
27. Smith KA, Conen F 2004 Impacts of land management on fluxes of trace greenhouse gases. *Soil Use Manage.* **20**, 255–263 (doi:10.1079/SUM2004238)
28. Yan X, Ohara T, Akimoto H 2003 Development of region-specific emission factors and estimation of methane emission from rice field in East, Southeast and South Asian countries. *Glob. Change Biol.* **9**, 237–254. (doi:10.1046/j.1365-2486.2003.00564.x)

*Phil. Trans. R. Soc. A.*

29. Cai ZC, Tsuruta H. & Minami, K. 2000 Methane emissions from rice fields in China: measurements and influencing factors. *J. Geophys. Res.* 105, 17 231–17 242. (doi:10.1029/2000JD900014)

30. Cai, Z. C., Tsuruta H, Gao M, Xu H, Wei CF. 2003 Options for mitigating methane emission from a permanently flooded rice field. *Glob. Change Biol.* 9, 37–45. (doi:10.1046/j.1365-2486.2003.00562.x)

31. Kang GD, Cai ZC, Feng, XZ 2002 Importance of water regime during the non-rice growing period in winter in regional variation of CH<sub>4</sub> emissions from rice fields during following rice growing period in China. *Nutr. Cycl. Agroecosyst.* 64, 95–100 (doi:10.1023/A:1021154932643)

32. Tian Z, Fan Y, Wang K, Zhong HL, Sun LX, Fan DL, Tubiello FN, Liu JG. 2021 Searching for “Win-Win” solutions for food-water-GHG emissions tradeoffs across irrigation regimes of paddy rice in China. *Resour Conserv Recycl.* 166, 105360, (doi: 10.1016/j.resconrec.2020.105360)

33. Xu H, Cai, ZC, Jia ZJ, Tsuruta H. 2000 Effect of land management in winter crop season on CH<sub>4</sub> emission during the following flooded and rice-growing period. *Nutr. Cycl. Agroecosyst.* 58, 327–332 (doi:10.1023/A:1009823425806)

34. Cai ZC, Xu H 2004 Options for mitigating CH<sub>4</sub> emissions from rice fields in China. In *Material circulation through agro-ecosystems in East Asia and assessment of its environmental impact* (ed. Y. Hayashi) NIAES Series, no. 5, pp. 45–55. Tsukuba, Japan: NIAES.

35. Blaxter KL, Clapperton JL 1965 Prediction of the amount of methane produced by ruminants. *Br. J. Nutr.* 19, 511–522 (doi:10.1079/BJN19650046)

36. Johnson KA, Johnson DE. 1995 Methane emissions from cattle. *J. Anim. Sci.* 73, 2483–2492 (doi:10.2527/1995.7382483x)

37. Lovett D, Lovell S, Stack L, Callan J, Finlay M, Connolly J, O’Mara, FP 2003 Effect of forage/concentrate ratio and dietary coconut oil level on methane output and performance of finishing beef heifers. *Livest. Prod. Sci.* 84, 135–146 (doi:10.1016/j.livprodsci.2003.09.010)

38. Beauchemin K, McGinn S 2005 Methane emissions from feedlot cattle fed barley or corn diets. *J. Anim. Sci.* 83, 653–661 (doi:10.2527/2005.833653x)

39. Phetteplace HW, Johnson DE, Seidl AF 2001 Greenhouse gas emissions from simulated beef and dairy livestock systems in the United States. *Nutr. Cycl. Agroecosyst.* 60, 9–102 (doi:10.1023/A:1012657230589)

- 
40. Lovett DK, Shalloo L, Dillon P, O'Mara FP 2006 A systems approach to quantify greenhouse gas fluxes from pastoral dairy production as affected by management regime. *Agric. Syst.* **88**, 156–179 (doi:10.1016/j.agsy.2005.03.006)
41. Machmülller A, Ossowski DA, Kreuzer M 2000 Comparative evaluation of the effects of coconut oil, oilseeds and crystalline fat on methane release, digestion and energy balance in lambs. *Anim. Feed Sci. Technol.* **85**, 41–60 (doi:10.1016/S0377-8401(00)00126-7)
42. Jordan E, Lovett DK, Hawkins M, O'Mara FP 2004 The effect of varying levels of coconut oil on methane output from continental cross beef heifers. In *Proc. Int. Conf. on Greenhouse Gas Emissions from Agriculture—Mitigation Options and Strategies* (ed. A Weiske), pp. 124–130. Leipzig, Germany: Institute for Energy and Environment
43. Leng RA 1991 *Improving ruminant production and reducing methane emissions from ruminants by strategic supplementation*. EPA report, no. 400/1-91/004. US Environmental Protection Agency, Washington, DC
44. McCrabb GJ, Kurihar M, Hunte, RA. 1998 The effect of finishing strategy of lifetime methane production for beef cattle in northern Australia. *Proc. Nutr. Soc. Aust.* **22**, 55
45. Alcock D, Hegarty RS 2005 Effects of pasture improvement on productivity, gross margin and methane emissions of grazing sheep enterprises. In *Second Int. Conf. on Greenhouse Gases and Animal Agriculture, Working Papers* (eds CR Soliva, J Takahashi, M Kreuzer), pp. 127–130. Zurich, Switzerland: ETH
46. Benz DA, Johnson DE 1982 The effect of monensin on energy partitioning by forage fed steers. *Proc. West Section Am. Soc. Anim. Sci.* **33**, 60–63
47. Van Nevel CJ, Demeyer DI. 1995 Lipolysis and biohydrogenation of soybean oil in the rumen in vitro: inhibition by antimicrobials. *J. Dairy Sci.* **78**, 2797–2806 (doi: 10.3168/jds.S0022-0302(95)76910-7)
48. McGinn SM, Beauchemin KA, Coates T, Colombatto D. 2004 Methane emissions from beef cattle: effects of monensin, sunflower oil, enzymes, yeast, and fumaric acid. *J. Anim. Sci.* **82**, 3346–3356 (doi:10.2527/2004.82113346x)

49. Rumpler WV, Johnson DE, Bates DB 1986 The effect of high dietary cation concentrations on methanogenesis by steers fed with or without ionophores. *J. Anim. Sci.* **62**, 1737–1741 (doi:10.2527/jas1986.6261737x)

50. Wolin EA, Wolf RS, Wolin MJ. 1964 Microbial formation of methane. *J. Bacteriol.* **87**, 993–998.

51. Van Nevel CJ, Demeyer DI. 1996 Influence of antibiotics and a deaminase inhibitor on volatile fatty acids and methane production from detergent washed hay and soluble starch by rumen microbes in vitro. *Anim. Feed Sci. Technol.* **37**, 21–31. (doi:10.1016/0377-8401(92)90117-O)

52. Newbold CJ, Rode LM 2005 Dietary additives to control methanogenesis in the rumen. In *Second Int. Conf. on Greenhouse Gases and Animal Agriculture, Working Papers* (eds CR Soliva, J Takahashi, M Kreuzer), pp. 60–70. Zurich, Switzerland: ETH

53. Newbold CJ, Ouda JO, Lopez S, Nelson N, Omed H, Wallace RJ, Moss AR 2002 Propionate precursors as possible alternative electron acceptors to methane in ruminal fermentation. In *Greenhouse gases and animal agriculture* (eds J Takahashi, BA Young), pp. 151–154. Amsterdam, The Netherlands: Elsevier

54. Newbold CJ, López S, Nelson N, Ouda JO, Wallace RJ, Moss AR. 2005 Propionate precursors and other metabolic intermediates as possible alternative electron acceptors to methanogenesis in ruminal fermentation in vitro. *Br. J. Nutr.* **94**, 27–35. (doi:10.1079/BJN20051445)

55. Wright ADG *et al.* 2004 Reducing methane emissions in sheep by immunization against rumen methanogens. *Vaccine* **22**, 3976–3985. (doi:10.1016/j.vaccine.2004.03.053)

56. Bauman DE. 1992 Bovine somatotropin: review of an emerging animal technology. *J. Dairy Sci.* **75**, 3432–3451 (doi:10.3168/jds.S0022-0302(92)78119-3)

57. Schmidely P 1993 Quantitative review on the use of anabolic hormones in ruminants for meat production. I. Animal performance. *Ann. Zootech.* **42**, 333–359.

58. Johnson DE, Ward GM, Torrent J. 1991 The environmental impact of bovine somatotropin (bST) use in dairy cattle. *J. Dairy Sci.* **74**, 209 (doi:10.2134/jeq1992.00472425002100020001x)

59. McCrabb GC 2001 Nutritional options for abatement of methane emissions from beef and dairy systems in Australia. In *Greenhouse gases and animal agriculture* (eds J Takahashi, BA Young), pp. 115–124. Amsterdam, The Netherlands: Elsevier.



60. Vijn S, Compart DP, Dutta N, Foukis A, Hess M, Hristov AN, Kalscheur KF, Kebreab E, Nuzhdin SV, Price NN, Sun Y, Tricarico JM, Turzillo A, Weisbjerg MR, Yarish C, Kurt TD. 2020 Key considerations for the use of seaweed to reduce enteric methane emissions from cattle. *Front. Vet. Sci.* **7**, 597430. (doi:10.3389/fvets.2020.597430)
61. Boadi D, Benchaar C, Chiquette J, Massé D. 2004 Mitigation strategies to reduce enteric methane emissions from dairy cows: update review. *Can. J. Anim. Sci.* **84**, 319–335 (doi:10.4141/A03-109)
62. Lovett DK, O'Mara FP 2002 Estimation of enteric methane emissions originating from the national livestock beef herd: a review of the IPCC default emission factors. *Tearmann* **2**, 77–83.
63. Clemens J, Ahlgrimm HJ. 2001 Greenhouse gases from animal husbandry: mitigation options. *Nutr. Cycl. Agroecosyst.* **60**, 287–300 (doi:10.1023/A:1012712532720)
64. Monteny GJ, Groenestein CM, Hilhorst MA. 2001 Interactions and coupling between emissions of methane and nitrous oxide from animal husbandry. *Nutr. Cycl. Agroecosyst.* **60**, 123–132 (doi:10.1023/A:1012602911339)
65. Monteny GJ, Bannink A, Chadwick D. 2006 Greenhouse gas abatement strategies for animal husbandry. *Agric. Ecosyst. Environ.* **112**, 163–170 (doi:10.1016/j.agee.2005.08.015)
66. Paustian K. et al. 2004. *Agricultural mitigation of greenhouse gases: science and policy options. Council on Agricultural Science and Technology (CAST) report*, R141 2004, ISBN 1-887383-26-3, 120pp, May 2004.
67. Clemens J, Trimborn M, Weiland P, Amon B 2006 Mitigation of greenhouse gas emissions by anaerobic digestion of cattle slurry. *Agric. Ecosyst. Environ.* **112**, 171–177 (doi:10.1016/j.agee.2005.08.016)
68. Gonzalez-Avalos E, Ruiz-Suarez LG 2001 Methane emission factors from cattle in Mexico. *Bioresour. Technol.* **80**, 63–71 (doi:10.1016/S0960-8524(01)00052-9)
69. Külling DR, Menzi H, Sutter F, Lischer P, Kreuzer M. 2003 Ammonia, nitrous oxide and methane emissions from differently stored dairy manure derived from grass- and hay-based rations. *Nutr. Cycl. Agroecosyst.* **65**, 13–22. (doi:10.1023/A:1021857122265)



- 
70. Pattey E, Trzcinski MK, Desjardins RL 2005 Quantifying the reduction of greenhouse gas emissions as a result of composting dairy and beef cattle manure. *Nutr. Cycl. Agroecosyst.* **72**, 173–187 (doi:10.1007/s10705-005-1268-5)
71. Bruun S, Stoumann Jensen L, Vu VTKV, Sommer S. 2014 Small-scale household biogas digesters: An option for global warming mitigation or a potential climate bomb? *Renew. Sust. Energ. Rev.* **33**, 736–741 (doi:10.1016/j.rser.2014.02.033)
72. Godoy JE, Camargo JR, Santos RR, et al. 2013 Design of biogas pipeline - energy and sanitation. Conference: 8th International Conference on Diffusion in Solids Liquids (DSL 2012). Istanbul, Turkey Jun 25-29, 2012. Diffusion in Solids and Liquids VIII Book Series: Defect and Diffusion Forum. 334–335. pp. 264-273
73. Smith KR, Uma R, Kishore VVN, Joshi KLV, Zhang J, Rasmussen RA, et al. 2000 Greenhouse gases from small-scale combustion devices in developing countries, Phase IIa. Household stoves in India. Washington, DC, USA: United States. Environmental Protection Agency. pp. 1–89.
74. Tumwesige V, Fulford D, Davidson GC. 2014 Biogas appliances in Sub-Sahara Africa. *Biomass Bioenerg.* **70**, 40-50 (doi:10.1016/j.biombioe.2014.02.017)
75. Bartoli A, Hamelin L, Rozakis S, Borzęcka M, Brandão M. 2019 Coupling economic and GHG emission accounting models to evaluate the sustainability of biogas policies. *Renew. Sust. Energ. Rev.* **106**, 133–148 (doi: 10.1016/j.rser.2019.02.031)
76. Subedi M, Matthews RB, Pogson M, Abegaz A, Balana BB, Oyesiku-Blakemore J, Smith J. 2014 Can biogas digesters help to reduce deforestation in Africa? *Biomass Bioenerg.* **70**, 87-98 (doi:10.1016/j.biombioe.2014.02.029)
77. Smith JU, Fischer A, Hallett PD, Homans HY, Smith P, Abdul-Salam Y, Emmerling HH, Phimister EC. 2015 Sustainable use of organic resources for bioenergy, food and water provision in rural Sub-Saharan Africa. *Renew. Sust. Energ. Rev.* **50**: 903-917 (doi:10.1016/j.rser.2015.04.071)
78. Smith J, Abegaz A, Matthews R, Subedi M, Orskov ER, Tumwesige V, et al. 2014 What is the potential for biogas digesters to improve soil carbon sequestration in sub-Saharan Africa? Comparison with other uses of organic residues, *Biomass Bioenerg.* **70**, 73-86 (doi:10.1016/j.biombioe.2014.01.056)
79. Bansal V, Tumwesige V, Smith JU. 2017 Water for small-scale biogas digesters in sub-Saharan Africa. *GCB. Bioenergy.* **9**(2), 339–357 (doi:10.1111/gcbb.12339)

- 
80. Wardle JM, Fischer A, Tesfaye Y, Smith J. 2021 Seasonal variability of resources: The unexplored adversary of biogas use in rural households in Ethiopia. *Current Res. Environ. Sust.* (submitted)
81. Khoiyangbam RS, Kumar S, Jain MC, Gupta N, Kumar A, Kumar V. 2004 Methane emission from fixed dome biogas plants in hilly and plain regions of northern India. *Bioresour. Technol.* **95**, 35–39 (doi:10.1016/j.biortech.2004.02.009)
82. Khoiyangbam RS, Kumar S, Jain MC. 2004 Methane losses from floating gasholder type biogas plants in relation to global warming. *J. Sci. Ind. Res.* **63**, 344–347 ([http://nopr.niscair.res.in/bitstream/123456789/5401/1/JSIR%2063\(4\)%20344-347.pdf](http://nopr.niscair.res.in/bitstream/123456789/5401/1/JSIR%2063(4)%20344-347.pdf))
83. Flesch TK, Desjardins RL, Worth D. 2011 Fugitive methane emissions from an agricultural biodigester. *Biomass Bioenerg.* **35**, 3927–3935 (doi:10.1016/j.biombioe.2011.06.009)
84. Prapasongsa T, Pholchan P, Hansen JA, Poulsen TG, Christensen P. 2009 Improved energy recovery efficiencies from piggery waste biogas plants in Thailand using Danish experiences. In: Proceedings of the world renewable energy congress, the 3rd international conference on sustainable energy and environment (SE2009, 18–23 May 2009, Bangkok, Thailand).
85. Porter SD, Reay DS, Higgins P, Bomberg E. 2016 A half-century of production-phase greenhouse gas emissions from food loss & waste in the global food supply chain. *Sci. Total Environ.* **571**, 721–729 (doi:10.1016/j.scitotenv.2016.07.041)
86. Ripple WJ, Smith P, Haberl H, Montzka SA, McAlpine C, Boucher DH 2014 Ruminants, climate change and climate policy. *Nat. Clim. Change* **4**, 2–5. (doi:10.1038/nclimate2081).
87. Tilman D, Clark M 2014 Global diets link environmental sustainability and human health. *Nature* **515**, 518–522 (doi:10.1038/nature13959)
88. Clark M, Tilman D 2017 Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. *Environ. Res. Lett.* **12**, 64016 (doi:10.1088/1748-9326/aa6cd5)
89. Poore J, Nemecek T. 2018 Reducing food's environmental impacts through producers and consumers. *Science* **360** 987–992 (doi:10.1126/science.aag0216)

- 
90. Springmann M *et al.* 2018 Options for keeping the food system within environmental limits. *Nature* **562**, 519–525 (doi:10.1038/s41586-018-0594-0)
91. Stehfest E, Bouwman L, Van Vuuren DP, Den Elzen MGJ, Eickhout B, Kabat P 2009 Climate benefits of changing diet. *Clim. Change* **95**, 83–102 (doi:10.1007/s10584-008-9534-6)
92. Popp A, Lotze-Campen H, Bodirsky B. 2010 Food consumption, diet shifts and associated non-CO<sub>2</sub> greenhouse gases from agricultural production. *Glob. Environ. Change* **20**, 451–462 (doi:10.1016/j.gloenvcha.2010.02.001)
93. Bajželj B, Richards KS, Allwood JM, Smith P, Dennis JS, Curmi E, Gilligan CA 2014 Importance of food-demand management for climate mitigation. *Nat. Clim. Change* **4**, 924–929 (doi:10.1038/nclimate2353)
94. Aleksandrowicz L, Green R, Joy EJM, Smith P, Haines A 2016 The impacts of dietary change on greenhouse gas emissions, land use, water use, and health: A systematic review. *PLoS ONE* **11**, e0165797 (doi:10.1371/journal.pone.0165797)
95. Roe S *et al.* 2019 Contribution of the land sector to a 1.5°C World. *Nat. Clim. Change* **9**, 817–828 (doi:10.1038/s41558-019-0591-9)
96. Quested T, Ingle R, Parry A 2013 Household food and drink waste in the United Kingdom 2012. WRAP, London.
97. UN FAO (Gerber P *et al.*) 2010 *Greenhouse Gas Emissions from the Dairy Sector. A Life Cycle Assessment*. UN Food and Agriculture Organization, Rome.
98. Thoma G *et al.* 2013 Greenhouse gas emissions from milk production and consumption in the United States: A cradle-to-grave life cycle assessment circa 2008. *Int. Dairy J.* **31**, S3-S14 (doi:10.1016/j.idairyj.2012.08.013).
99. Reay D. 2019 Climate-Smart Milk. In *Climate-Smart Food* (Reay D) pp. 49-66). Palgrave Pivot, Cham.
100. Zhang B *et al.* 2016 Methane emissions from global rice fields: Magnitude, spatiotemporal patterns, and environmental controls. *Global Biogeochem. Cycles* **30**, 1246-1263 (doi: 10.1002/2016GB005381)
101. Wang C *et al.* 2018 An additive effect of elevated atmospheric CO<sub>2</sub> and rising temperature on methane emissions related to methanogenic community in rice paddies. *Agric. Ecosyst. Environ.* **257**, 165-174 (doi:10.1016/j.agee.2018.02.003)

- 
102. Bryngelsson D, Hedenus F, Johansson DJA, Azar C, Wirsenius S. 2017 How do dietary choices influence the energy-system cost of stabilizing the climate? *Energies* **10**, 182. (doi:10.3390/en10020182)
103. Lynch J, Cain M, Frame D, Pierrehumbert R. 2021 Agriculture's contribution to climate change and role in mitigation is distinct from predominantly fossil CO<sub>2</sub>-emitting sectors. *Front. Sustain. Food Syst.* **4**, 518039. (doi: 10.3389/fsufs.2020.518039)
104. Leahy S, Clark H, Reisinger A. 2020 challenges and prospects for agricultural greenhouse gas mitigation pathways consistent with the Paris Agreement. *Front. Sustain. Food Syst.* **4**, 69. (doi: 10.3389/fsufs.2020.00069)
105. Hawken P (Ed.) 2017 *Drawdown: The most comprehensive plan ever proposed to reverse global warming*. 240pp, New York, NY: Penguin.
106. IPCC. 2018 Summary for Policymakers. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty* [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)]. World Meteorological Organization, Geneva, Switzerland, 32 pp.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


---

**Figure captions**

**Figure 1.** Global methane emissions (Tg y<sup>-1</sup>) from global emissions sectors in 2017. Bottom up best estimates from [11].

**Figure 2.** Agricultural methane emissions 1961-2017 by source. Data from [1]. Note: Though savannas are used to varying extents for grazing domestic livestock, savanna burning emissions are not included.

**Figure 3.** Estimated maximum mitigation potential (for carbon price of 100 US\$ tCO<sub>2</sub>e<sup>-1</sup>) for methane emissions from agriculture [10,23]. Note: Biogas mitigation potential not shown as emission reductions are accounted for in the energy sector.



■ Wetlands                      ■ Other natural sources                      ■ Agriculture  
■ Waste                      ■ Fossil fuels                      ■ Biomass and biofuel burning

